

Antibody Datasheet / Certificate of Analysis

Product Name:	Mouse anti-Zika virus NS1
Clone number:	D11
Isotype:	Mouse IgG1
Product code:	AbZIKVNS1-D11-100
Batch Number:	16070800
Immunogen:	Full length recombinant NS1 protein of Zika virus produced in HEK293 cells (available from the Native Antigen Company, product code ZIKV-NS1-100)
Amount:	100ug
Concentration:	1.0 mg/ml
Buffer:	Phosphate Buffered Saline pH7.4
Preservative:	None present. 0.2um filtered.
Fusion partners:	Spleen cells from immunised Balb/c mice were fused with cells from the SP2/0-Ag14 myeloma cell line.
Purification:	Antibody was purified from hybridoma cell culture supernatant by affinity chromatography on Protein G
Specificity:	This antibody is specific for the NS1 protein of Zika virus, detecting NS1 from both the Uganda and Suriname strains. It demonstrates negligible cross-reactivity with NS1 proteins from Dengue virus (all serotypes), Japanese Encephalitis Virus and Yellow Fever Virus. A small amount of cross-reactivity has been observed with the NS1 protein from West Nile Virus in a direct ELISA (see results below and Images).

Applications: Direct ELISA (NS1 antigen bound to plate)

Sandwich ELISA (as both capture and detection antibody). This antibody may be used in combination with clone B4 (product code AbZIKVNS1-B4)

This antibody is not recommended for use in Western Blot

This antibody is suitable for use in immunofluorescence applications (see results below and [Images](#))

Antigen background: Zika virus is an emerging disease that is spread by *Aedes* mosquitoes. The virus was first isolated in Central Africa, and has since spread to South Asia and more recently to South America. It is a member of the flavivirus family, and is structurally closely related to viruses such as Dengue Fever Virus. Outbreaks were reported in Micronesia in 2007 and in Brazil in 2015, confirming at least 13 autochthonous infections. The Zika virus outbreak in Brazil in 2016 has gained world-wide attention, and has been linked to an increasing number of microcephaly cases. In April 2016 the Centers for Disease Control, in the USA, confirmed the link between Zika virus infection of the fetus with microcephaly.


Clinically Zika virus can cause mild fever, rash, myalgia, arthralgia and headaches, with one in four infected individuals being asymptomatic. Due to similar symptoms Zika virus infected individuals can easily be mis-diagnosed as a dengue infection and vice-versa. In addition, Zika virus has been implicated in causing microcephaly through transmission *in utero*. There is no vaccine or specific treatment available for Zika virus.

The NS1 protein is a major non-structural protein expressed by the Zika Virus. The NS1 monomer is a glycosylated protein of approximately 45kD, which associates with lipids and forms a homodimer inside infected cells. It is necessary for viral replication, and is also secreted into the extracellular space as a hexameric lipoprotein particle, which is involved in immune evasion and pathogenesis by interacting with components from both the innate and adaptive immune systems, as well as other host factors. NS1 is one of the major antigenic markers for viral infection with Zika.

Storage: Store at +4°C for up to one week, or at -20°C for longer periods
For long term storage at +4°C the addition of 0.09% w/v sodium azide is recommended.
The antibody is shipped at ambient temperature.
Avoid repeated freeze/thaw cycles.


Direct ELISA data

An ELISA plate was coated with 100ng of antigen per well, then blocked with 2% BSA. Primary antibody was used at a concentration of 1ug/ml, and the detection antibody used was Goat anti-mouse IgG:HRP (Bio-Rad, 1:2000). The substrate used was TMB (KPL).




Sandwich Assay Data

An ELISA plate was coated with 500ng of coating antibody, then blocked with Casein, and samples were added at a range of concentrations. Detection antibody was biotinylated, and used at a concentration of 0.25ug/ml. Visualisation was with Streptavidin:HRP and the substrate used was TMB (KPL).



Comparison of sandwich assay antibody variations



Immunofluorescence data

Vero cells were seeded on coverslips infected with ZIKV (African or Asian strains) for 30h at MOI 0.5. Control coverslips consisted of uninfected Vero cells. After fixation with 4% PFA, samples were stained with Mouse anti-Zika virus NS1 (D11). Antibody was diluted 1:500 and Triton X-100 was used as detergent. Imaging was performed using a Leica SP5 confocal microscope. Immunofluorescence for both African and Asian strains was observed; the African strain illustrated below (Virology Research Services).

