


Antibody Datasheet

| | |
|-------------------------------|--|
| Product Name: | Rabbit anti <i>Borrelia burgdorferi</i> ErpN/OspC |
| Product: | Purified rabbit anti ErpN/OspC antibody, unconjugated |
| Product Type: | Polyclonal |
| Isotype: | Rabbit IgG |
| Product code: | PAB21452-100 |
| Batch Number: | R001218 |
| Amount: | 0.1 ml (1.0 mg/mL by UV absorbance at 280 nm) |
| Physical State: | Lyophilized |
| Buffer: | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Reconstitution Vol: | 100 µL |
| Reconstitution Buffer: | Deionized water (or equivalent) |
| Preservative: | 0.01% (w/v) Sodium Azide |
| Immunogen: | Recombinant MBP tagged <i>B. burgdorferi</i> ErpN/OspC protein |
| Purification: | Protein-A purified and cross-adsorbed against MBP from monospecific antiserum by chromatography |
| Specificity: | This antibody is directed against, and shows specific reactivity for <i>Borrelia burgdorferi</i> OspC protein. Reactivity with ErpN/OspC protein from other sources has not been determined. |
| Applications: | ELISA (1:5000), WB (1:1000) |



Storage:

Store at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. Antibody is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.



Western blot showing detection of 0.1 µg recombinant proteins in Western blot. Lane 1: Molecular weight markers. Lane 2: MBP-ErpN/OspE fusion protein (arrow; 59.5 kDa expected MW). Lane 3: fusion protein (MBP-tagged) plus cleaved fusion proteins (without MBP). Lane 4: MBP alone. The lower bands are probably breakdown products. The upper bands in lane 3 are fusion protein (top band), or breakdown products of the fusion protein (bands in middle of blot). Protein was run on a 4-20% gel, then transferred to 0.45 µm nitrocellulose. After blocking with 1% BSA-TTBS overnight at 4°C, primary antibody was used at 1:1000 at room temperature for 30 min. HRP-conjugated Goat-Anti-Rabbit secondary antibody was used at 1:40,000 in blocking buffer and imaged on the VersaDoc™ MP 4000 imaging system (Bio-Rad).

