

Langford Locks Kidlington Oxfordshire OX5 1LH

Telephone: +44(0)1865 595230

Antibody Datasheet

Name: Anti-Human IgA + IgG + IgM (H+L) Antibody, Mouse Serum Adsorbed

Product Code: 5210-0164

Batch #:

Date of Manufacture:

Product Description: This broad spectrum reagent contains a mixture of 3 antibodies that recognize: human IgA, IgG and IgM. Affinity purified antibody isolated from a pool of serum from goats immunized with purified human IgA, IgG, and IgM. Each lot is tested to assure specificity and lot-to-lot consistency using an in-house ELISA assay.

Amount: 1.0 mg

Presentation: Lyophilized

Rehydration: Rehydration of antibodies in TBS or buffers other than those listed here is not

recommended.

Procedure A: 50% Glycerol

At a working dilution, the level of glycerol is too small to affect most assays. The use of glycerol is not recommended when the antibody is used in live cell work. Rehydration - Add 1 mL of 50% glycerol in water to the vial. Pipette up and down several times to ensure proper mixing. Storage: This product may be stored either refrigerated or frozen as desired in 50% glycerol. Stable for a minimum of 1 year. Storage: This product may be stored either refrigerated or frozen as desired in 50% glycerol. Stable for a minimum of 1 year.

Procedure B: 2mM Acetic Acid Solution

Rehydration - Rehydrate with 1 mL of 2mM Glacial Acetic acid made with reagent quality water. Rotate the vial until the lyophilized pellet is totally dissolved. Storage: This product may be stored for up to 1 week refrigerated; thereafter, it should be stored frozen. Stable for a minimum of 1 year at -20°C.

Stabilizer: No stabilizers added

Preservative: No preservatives added. Non-sterile.

Purification: Affinity purified

Specificity: Tested by gel diffusion and ELISA techniques as applicable. This product reacts specifically with human IgA, IgG, and IgM and may recognize other immunoglobulin types that have light chains in common with IgA, IgG, and IgM.







Usage Guidelines

Therapeutic Use.

Langford Locks Kidlington Oxfordshire OX5 1LH

Telephone: +44(0)1865 595230

Cross-Reactivity: Antibodies to human IgA, IgG, and IgM may cross-react with immunoglobulins of other mammalian species if common binding sites are shared. Cross-reactivity with mouse serum has been minimized with affinity procedures.

Storage:	Store at +4°C until rehydrated. Once rehydrated store at -20°C
Χ	
QC	
X	
QA	

Products are for Research Use or for Further Manufacturing Use only. Not for Diagnostic or



